

ϵ -Expansion of the Conductivity at the Superconductor-Mott Insulator Transition

Rosario Fazio⁽¹⁾ and Dario Zappalà⁽²⁾

⁽¹⁾*Institut für Theoretische Festkörperphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany**

⁽²⁾*Dipartimento di Fisica, Università di Catania, and INFN, sezione di Catania, corso Italia 57, 95128 Catania, Italy*

Abstract

We study the critical behavior of the conductivity $\sigma(\omega)$ at the zero temperature superconductor-Mott insulator transition in d space-time dimensions for a model of bosons with short-range interaction and no disorder. We obtain $\sigma(\omega_n) = (4e^2/\hbar)\sigma_\epsilon\omega_n^{1-\epsilon}$, as predicted by the scaling theory, and the prefactor σ_ϵ is calculated in the ϵ -expansion, to order ϵ^2 ($\epsilon = 4 - d$). In two spatial dimensions, ($d = 3$), we find a value of the universal conductance $\sigma^* = 0.315(4e^2/h)$, in good agreement with the known Monte Carlo results.

PACS numbers: 74.75 +t, 74.40 +k

In two dimensions at the $T = 0$ superconductor-insulator (SI) transition the conductance is finite and universal^{1,2}. Since the original prediction of a metallic behavior at zero temperature for two dimensional superconductors there has been a lot of interest both on experimental and theoretical side.

The systems where the $T = 0$ SI transition has been observed are ultra thin superconducting films³ and Josephson junction arrays⁴⁻⁶. The films are driven through the SI transition by varying the thickness while, for the arrays, the ratio of the charging energy to the Josephson coupling determines whether they are in the insulating or in the superconducting phase.

Many theoretical works have investigated various aspects of the SI transition. The superconductor-Mott insulator transition and the universal conductance in a model with no disorder was considered in Ref. 7 by means of $1/N$ expansion and Monte Carlo simulations. The duality between charges and vortices, leading to an universal conductance of $(4e^2/h)^7$, has been discussed in Ref. 8. Fisher *et al.*⁹ studied the disordered interacting boson problem using a scaling approach. The dirty boson system and the transition to the Bose glass phase (including the case of long-range Coulomb interaction) was extensively studied in^{10,11}. Wen employed a scaling theory of conserved current at anisotropic critical points¹² identifying many universal amplitudes. One of these amplitudes in two dimensions reduces to the universal conductance σ^* . The finite frequencies properties close to the transition point were analyzed by means of the $1/N$ expansion^{13,14}.

The determination of the value of σ^* relies almost entirely on numerical methods (Monte Carlo¹⁵ and exact diagonalization¹⁶). The only analytical calculation of σ^* is based on the $1/N$ expansion⁷. Another powerful and widely used method to calculate the critical properties is the ϵ -expansion (for a review see Ref. 17).

In this Letter the conductivity at the SI transition is calculated applying the ϵ -expansion. We will compute the scaling form of the $\sigma(\omega)$ discussed in Refs. 1,12 and we will determine, in two spatial dimensions, the value of σ^* .

The description of the SI transition starts from the hypothesis that the critical properties are determined by the quantum fluctuations of the phase of the superconducting order parameter. A widely used model for the SI transition is a Hubbard like model for strongly interacting bosons on a lattice (Bose-Hubbard model)^{9,10,18}.

$$H = \frac{U}{2} \sum_i n_i(n_i - 1) - \mu \sum_i n_i - \frac{t}{2} \sum_{\langle ij \rangle} (b_i^\dagger b_j + h.c.) \quad (1)$$

where $b^\dagger, b$ are the creation and annihilation operators for bosons and n_i is the number of bosons. U describes the short range interaction between bosons, μ is the chemical potential and t is the hopping matrix element. At integer fillings, the Bose-Hubbard model in $d - 1$ spatial dimensions is in the same universality class of the XY model in d dimensions. It is more convenient for computational purposes to work with the Ginzburg-Landau action (onto which the Bose-Hubbard model maps)

$$F[\psi] = \int d^{d-1} \vec{r} d\tau \left\{ \delta_0 |\psi|^2 + |\nabla \psi|^2 + |\partial_\tau \psi|^2 + \frac{\kappa^\epsilon u_0}{2} |\psi|^4 \right\}. \quad (2)$$

where $\psi(\vec{r}, \tau)$ is a complex one-component scalar field. In (2) we have introduced $\epsilon = 4 - d$ and the arbitrary momentum scale κ in order to keep the quartic coupling u_0 dimensionless.

The bare parameters δ_0 and u_0 can be related to the coupling constants in the Bose-Hubbard model^{9,13}. u_0 includes the effect of the interaction and δ_0 measures of the distance from the critical point (we set $\delta_0 = 0$ in the following). It is evident from eq. (2) that the quantum system at zero temperature is equivalent to a classical system in d dimensions.

The conductivity of the system can be obtained by minimally coupling the scalar field in eq. (2) to a vector potential $\vec{A}$. The imaginary time frequency-dependent conductivity is given by

$$\sigma_{\mu\nu}(\omega_n) = \frac{\hbar}{\omega_n} \int d^{d-1}r d\tau \frac{\delta^2 \ln Z}{\delta A_\mu(r, \tau) \delta A_\nu(0)} \Big|_{\vec{A}=0} e^{i\omega_n \tau} \quad (3)$$

where Z is the partition function of the system. The conductivity is obtained by analytically continuing eq. (3) to real frequencies ($i\omega_n = \omega + i0^+$). Being interested in the zero field case, we will only consider the longitudinal conductance (e.g. $\sigma_{xx}(\omega_n)$), which we shall generically indicate as $\sigma(\omega_n)$.

By performing the functional derivatives it is straightforward to express the conductivity in terms of two and four point Green's functions. Following Cha et al.⁷ it is convenient to introduce the notation $\sigma(\omega_n) = (4e^2/\hbar)\rho(\omega_n)/\omega_n$ where

$$\rho(\omega_n) = 2 \int \frac{d^d q}{(2\pi)^d} \langle \psi_{\vec{q}}^* \psi_{\vec{q}} \rangle - 4 \int \frac{d^d q}{(2\pi)^d} \frac{d^d p}{(2\pi)^d} q_x p_x \langle \psi_{\vec{q}-\frac{\vec{k}}{2}}^* \psi_{\vec{p}+\frac{\vec{k}}{2}}^* \psi_{\vec{p}-\frac{\vec{k}}{2}} \psi_{\vec{q}+\frac{\vec{k}}{2}} \rangle. \quad (4)$$

In eq. (4) the $\vec{q}$ are real vectors in the d -dimensional space and $\vec{k} = (0, 0, \dots, \omega_n)$.

The two-point function in eq. (4), expressed in terms of the self-energy $\Sigma(\vec{q})$ is (in the following we shall use the notation $q = |\vec{q}|$): $\langle \psi_{\vec{q}}^* \psi_{\vec{q}} \rangle = G(\vec{q}) = (q^2 + \Sigma(\vec{q}))^{-1}$. Since the three point irreducible vertex is zero in the insulating phase where no spontaneous symmetry breaking occurs, the four point function in eq. (4) is the sum of a connected part containing the irreducible four point vertex $\Gamma^{(4)}(\vec{q}, \vec{p}, \vec{k})$, and two non-connected parts of which only one gives non vanishing contribution to $\rho(\omega_n)$. $\Gamma^{(4)}(\vec{q}, \vec{p}, 0)$ and $\Sigma(\vec{q})$ are related by the following Ward identity

$$\frac{\partial}{\partial q_x} \Sigma(\vec{q}) = 2 \int \frac{d^d q}{(2\pi)^d} p_x G^2(\vec{p}) \Gamma^{(4)}(\vec{q}, \vec{p}, 0). \quad (5)$$

This identity, related to the underlying gauge symmetry of the free energy in eq.(2), is essential to prove that $\rho(\omega_n = 0) = 0$. This can be easily checked by evaluating the derivative of the two-point Green's function and substituting in eq. (4). In the superfluid phase the cancelation does not occur due to the presence of three point vertex in the theory.

In order to get a more suitable expression of $\rho(\omega_n)$ for a perturbative analysis, we split the four point irreducible vertex in five parts, $\Gamma^{(4)}(\vec{q}, \vec{p}, \vec{k}) = \Gamma_o^{(4)} + \Gamma_s^{(4)}(\vec{q} + \vec{p}) + \Gamma_t^{(4)}(\vec{q} - \vec{p}) + \Gamma_u^{(4)}(\vec{k}) + \Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k})$ according to the external momenta dependence. $\Gamma_o^{(4)}$ has no momentum dependence and $\Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k})$ is the residual part with momentum dependence different from $\vec{q} + \vec{p}$ or $\vec{q} - \vec{p}$ or $\vec{k}$. This splitting turns out to be interesting because $\Gamma_o^{(4)}$ and $\Gamma_u^{(4)}(\vec{k})$, when inserted in eq. (4), vanish and $\Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k})$ gives a contribution to eq. (4), in the ϵ -expansion, only of order $O(\epsilon^3)$. Eq.(5) and the decomposition of the $\Gamma^{(4)}$ allow, after some algebra, to express $\rho(\omega_n)$ in the form

$$\begin{aligned}
\frac{\rho(\omega_n)}{4} &= \int \frac{d^d q}{(2\pi)^d} \left[q_x^2 + q_x \frac{\partial}{\partial q_x} \Sigma(\vec{q}) \right] G(\vec{q}) [G(\vec{q}) - G(\vec{q} + \vec{k})] \\
&- \int \frac{d^d q}{(2\pi)^d} \frac{d^d p}{(2\pi)^d} q_x p_x G(\vec{q}) G(\vec{p}) \left\{ [G(\vec{q} + \vec{k}) - G(\vec{q})] \gamma^{(4)}(\vec{q} - \vec{p}) [G(\vec{p} + \vec{k}) - G(\vec{p})] \right. \\
&\left. - G(\vec{p}) [2 G(\vec{q} + \vec{k}) - G(\vec{q})] \Gamma_{res}^{(4)}(\vec{q}, \vec{p}, 0) + G(\vec{p} + \vec{k}) G(\vec{q} + \vec{k}) \Gamma_{res}^{(4)}(\vec{q} + \frac{\vec{k}}{2}, \vec{p} + \frac{\vec{k}}{2}, \vec{k}) \right\} \quad (6)
\end{aligned}$$

where we have introduced $\gamma^{(4)}(\vec{q} - \vec{p}) = \Gamma_t^{(4)}(\vec{q} - \vec{p}) - \Gamma_s^{(4)}(\vec{q} - \vec{p})$. So far no approximations have been used to obtain eq. (6). The next step is the computation of the self-energy and four point vertex in the framework of the ϵ -expansion.

At the critical point $G^{-1}(\vec{q}) = q^{2-\eta}$ and the value of the critical exponent η , for the theory considered, is known to be $\eta = (\epsilon^2/50)(1 + 19\epsilon/20) + O(\epsilon^4)$. The expansion of $G(\vec{q})$ in powers of ϵ follows straightforwardly.

Let us now consider the four point vertex $\Gamma^{(4)}(\vec{q}, \vec{p}, \vec{k})$; its expansion in terms of the dimensionless renormalized coupling constant u can be formally written as

$$\Gamma^{(4)}(\vec{q}, \vec{p}, \vec{k}) = \kappa^\epsilon \left[-2u + u^2 A(\vec{q}, \vec{p}, \vec{k}) + u^3 B(\vec{q}, \vec{p}, \vec{k}) + O(u^4) \right] \quad (7)$$

It is understood that the function in eq. (7) is renormalized and therefore, employing dimensional regularization, no pole at $\epsilon = 0$ is present in $A(\vec{q}, \vec{p}, \vec{k})$ and $B(\vec{q}, \vec{p}, \vec{k})$. In the following calculation the minimal subtraction renormalization prescription has been used.

Since we are concerned with the computation of the four point vertex at the transition, we must evaluate eq. (7) at $u = u^*$, u^* being the fixed point of the β function of the theory: $\beta(u^*) = 0$. By introducing the counterterms a_i through the expansion of the bare coupling u_0 appearing in eq. (2): $u_0 = u(1 + a_1 u + a_2 u^2 + \dots)$, we find from the evaluation of the bare four point vertex at two loop in the minimal subtraction scheme: $a_1 = 5/(8\pi^2\epsilon)$, $a_2 = (25/\epsilon^2 - 15/(2\epsilon))/(8\pi^2)^2$.

The fixed point u^* , obtained as a function of the counterterms a_i (see for instance Ref. 19), is $u^* = (8\pi^2)(\epsilon/5 + 3\epsilon^2/25) + O(\epsilon^3)$. Since the coefficients $A(\vec{q}, \vec{p}, \vec{k})$ and $B(\vec{q}, \vec{p}, \vec{k})$ do not contain any pole at $\epsilon = 0$, a full $O(\epsilon^2)$ calculation of the four point vertex requires the computation of the one loop part $A(\vec{q}, \vec{p}, \vec{k})$ only.

Nevertheless, let us consider the momentum dependent structure of both $A(\vec{q}, \vec{p}, \vec{k})$ and $B(\vec{q}, \vec{p}, \vec{k})$. In the one loop part $A(\vec{q}, \vec{p}, \vec{k})$ one has terms like $\epsilon^n \log^{n+1}$, $\epsilon^{n+1} \log^{n+1}$, $\epsilon^{n+2} \log^{n+1}$, ..., with $n = 0, 1, 2, \dots$ and $\log$ being the logarithm of a generic combination of external momenta; in the two loop part $B(\vec{q}, \vec{p}, \vec{k})$ one gets $\epsilon^n \log^{n+2}$, $\epsilon^n \log^{n+1}$, $\epsilon^{n+1} \log^{n+1}$, As mentioned above, a full $O(\epsilon^2)$ computation of eq. (7) requires just the coefficient of the term $\epsilon^0 \log$ in $A(\vec{q}, \vec{p}, \vec{k})$, but, since we are interested in summing formally the leading $\log$ series in eq. (7), we need to know at least the first two terms of the series. This amounts to calculate the terms $\epsilon \log^2$ in $A(\vec{q}, \vec{p}, \vec{k})$ and $\epsilon^0 \log^2$ in $B(\vec{q}, \vec{p}, \vec{k})$.

In the minimal subtraction scheme $A(\vec{q}, \vec{p}, \vec{k})$ is

$$\begin{aligned}
A(\vec{q}, \vec{p}, \vec{k}) &= \frac{2}{(8\pi^2)} \left[C - \log \frac{|\vec{q} + \vec{p}|}{\kappa} - 2 \log \frac{|\vec{q} - \vec{p}|}{\kappa} - 2 \log \frac{|\vec{k}|}{\kappa} \right. \\
&\left. + \frac{\epsilon}{2} \log^2 \frac{|\vec{q} + \vec{p}|}{\kappa} + \epsilon \log^2 \frac{|\vec{q} - \vec{p}|}{\kappa} + \epsilon \log^2 \frac{|\vec{k}|}{\kappa} + O(\epsilon) \right] \quad (8)
\end{aligned}$$

where, according to our task, we have displayed all the contributions which are relevant for the computation of the first two terms of the leading $\log$ series and neglected the subleading logarithms ($\epsilon \log$) and the higher powers of ϵ . C in eq. (8) is a $O(1)$ number, irrelevant for the determination of $\rho(\omega_n)$ in eq. (6).

Still we need the $\epsilon^0 \log^2$ part of $B(\vec{q}, \vec{p}, \vec{k})$, which we indicate as $\hat{B}(\vec{q}, \vec{p}, \vec{k})$; it is

$$\hat{B}(\vec{q}, \vec{p}, \vec{k}) = -\frac{10}{(8\pi^2)^2} \left(\log^2 \frac{|\vec{q} + \vec{p}|}{\kappa} + 2 \log^2 \frac{|\vec{q} - \vec{p}|}{\kappa} + 2 \log^2 \frac{|\vec{k}|}{\kappa} \right) \quad (9)$$

As stated above, to order ϵ^2 , $\Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k}) = 0$, making eq. (6) much simpler. Actually the first nonvanishing contribution to $\Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k})$ comes from the subleading term $\epsilon^0 \log$ in $B(\vec{q}, \vec{p}, \vec{k})$ which is neglected in our calculation. We are now able to sum formally the leading $\log$ series obtaining some power of the momenta involved (note that the momentum-independent term $O(\epsilon)$ in the series is irrelevant because it vanishes when integrated in eq. (7)). Putting together eqs.(7, 8, 9), $\gamma^{(4)}(\vec{q} - \vec{p})$ becomes

$$\gamma^{(4)}(\vec{q} - \vec{p}) = -\frac{16\pi^2\epsilon}{25} |\vec{q} - \vec{p}|^\epsilon + O(\epsilon^3) \quad (10)$$

The importance of summing the leading $\log$ series is now clear: eq. (10) is correct only to order $O(\epsilon^2)$, but the infrared singularities due to the logarithms in eqs. (8, 9), disappear in the resummed quantity $|\vec{q} - \vec{p}|^\epsilon$.

Another comment is in order: the last result does not depend on the arbitrary momentum scale κ introduced in eq. (2), nor on the choice of the minimal subtraction scheme. Indeed, one can easily prove that the shift of the counterterms a_i by a finite (no poles at $\epsilon = 0$) amount, $a_i \rightarrow a_i + \bar{a}_i$, does not change $\gamma^{(4)}(\vec{q} - \vec{p})$ in eq. (10).

Finally, going back to eq. (6), we can insert eq. (10), and discard the terms containing $\Gamma_{res}^{(4)}(\vec{q}, \vec{p}, \vec{k})$, thus obtaining $\rho(\omega_n)$ and, consequently, $\sigma(\omega_n)$ to order ϵ^2 ; in dimensions lower than 4 all integrals in eq. (6) are convergent in the ultraviolet (in $d \geq 4$, a short-range cutoff should be introduced to regularize the results, but this is not important for our purposes) and we find

$$\sigma(\omega_n) = \frac{4e^2}{\hbar} \left\{ \left(\frac{\epsilon^2}{25} - 2 \right) \frac{\Gamma^2(d/2)\Gamma(1-d/2)}{\pi^{d/2} 2^d \Gamma(d)} - 2 z(d) \int \frac{d^d y}{y^4} f_{\vec{k}}(\vec{y}) f_{\vec{k}}^*(\vec{y}) \right\} \omega_n^{1-\epsilon} \quad (11)$$

where

$$f_{\vec{k}}(\vec{y}) = \int_0^1 d\alpha \int_0^\infty \frac{d\lambda}{(4\pi)^{d/2}} \frac{y_x}{2 \lambda^{d/2}} \exp\left(-\frac{y^2}{4\lambda}\right) \left[1 - \exp(-\alpha(1-\alpha)\lambda - i\alpha y_z) \right] \quad (12)$$

and $z(d) = \left[\pi^{d/2} / (\Gamma(d/2)) \int_0^\infty dx x^{d-5} \left(1 - (x/2)^{1-d/2} \Gamma(d/2) J_{(d/2-1)}(x) \right) \right]^{-1}$. $\Gamma(x)$ and $J_\nu(x)$ are, respectively, the Gamma and Bessel functions and the ω_n independent factor in eq. (11) has been expressed in integral form for the sake of simplicity.

Eq.(11) is the central result of our paper. The form of the frequency dependent conductivity, as predicted from the scaling theory^{1,12}, is obtained.

In the calculation of the $\sigma(\omega_n)$, ϵ -expansion was used to determine the irreducible vertices. The d -dimensional integrals in eq.(6) have been instead performed not resorting to any

expansion. This observation is crucial in order not to obtain spurious dependences on the ultraviolet cut-off needed to regularize the theory in higher dimensions. In this sense the calculation of the conductivity to $O(\epsilon^2)$ should be understood as the scheme of approximation for the vertices.

In two spatial dimension the universal conductance σ^* can be evaluated from eq. (11) at $d = 3$; this leads to the result displayed in Table 1, together with the other known determinations of σ^* . Our estimate is in agreement with the results of the Monte Carlo simulations in Ref. 7 within 10% but it should be noted that the $1/N$ and ϵ -expansion approach the Monte Carlo value from different sides.

As a final remark we want to go back to the connection with the experiments. The model with only phase fluctuations is certainly applicable to Josephson junction arrays, but it has been recently questioned in the case of thin films. New measurements²⁰ point out that the value of the conductance may be not universal and this should be associated to the relevance of fermionic degrees of freedom at the transition. This is also the outcome of recent calculations which include fermions starting from a Bose model with local damping²¹ or simulating a model of electrons interacting through an attractive potential²².

ACKNOWLEDGMENTS

We would like to thank A. Bonanno, V. Branchina, C. Bruder, A. van Otterlo, G. Schön, K-H. Wagenblast, and G.T. Zimanyi for useful discussions. The financial support of the European Community (HCM-network CHRX-CT93-0136), the Deutsche Forschungsgemeinschaft through SFB 195 and INFN is gratefully acknowledged.

REFERENCES

- * Permanent address. Istituto di Fisica, Università di Catania, viale A. Doria 6, 95129 Catania, Italy
- ¹ M.P.A. Fisher, G. Grinstein and S.M. Girvin, Phys. Rev. Lett. **64**, (1990) 587.
- ² X.G. Wen and A. Zee, Int.J.Mod.Phys. **B4**, 437 (1990).
- ³ D.B. Haviland, Y. Liu, and A.M. Goldman, Phys. Rev. Lett. **62**, 2180 (1989).
- ⁴ L.J. Geerligs, M. Peters, L.E.M. de Groot, A. Verbruggen, and J.E. Mooij, Phys. Rev. Lett. **63**, 326 (1989).
- ⁵ H.S.J van der Zant, L.J. Geerligs, and J.E. Mooij, Europhys. Lett. **19**, 541 (1992).
- ⁶ C.D. Chen, P. Delsing, D.B. Haviland, Y. Harada, and T. Claeson, Phys. Rev. **B50**, 3959 (1995).
- ⁷ M.-C. Cha, M.P.A. Fisher, S.M. Girvin, M. Wallin, A.P. Young, Phys. Rev. B **44**, (1991) 6883.
- ⁸ R. Fazio and G. Schön, Phys. Rev. B **43**, 5307 (1991).
- ⁹ M.P.A. Fisher, B. P. Weichman, G. Grinstein, and D. S. Fisher, Phys. Rev. B **40**, 546 (1989).
- ¹⁰ G. G. Batrouni, R. T. Scalettar, and G. T. Zimanyi, Phys. Rev. Lett. **65**, 1765 (1990); W. Krauth and N. Trivedi, Europhys. Lett. **14**, 627 (1991).
- ¹¹ E.S. Sørensen, M. Wallin, S.M. Girvin, and A.P. Young, Phys. Rev. Lett. **69**, 828 (1992).
- ¹² X.G. Wen, Phys. Rev. **B46**, 2655 (1992).
- ¹³ A. van Otterlo, K.-H. Wagenblast, R. Fazio, and G. Schön, Phys. Rev. **48** (1993) 3316.
- ¹⁴ A.P. Kampf and G. T. Zimanyi, Phys. Rev. **47** (1993) 279.
- ¹⁵ M. Wallin, E.S. Sørensen, S.M. Girvin, and A.P. Young, Phys. Rev. **B45**, 13136 (1992).
- ¹⁶ K. Runge, Phys. Rev. **B49**, 12115 (1994).
- ¹⁷ V. Privman, P.C. Hohenberg, and A. Aharony in *Phase Transitions and Critical Phenomena*, Vol.14 pag.1, Edited by C. Domb and J.L.L. Lebowitz, (Academic Press, London)
- ¹⁸ C. Bruder, R. Fazio, and G. Schön, Phys. Rev. B **47**, 342 (1993);
- ¹⁹ D. J. Amit *Field theory, the renormalization group, and critical phenomena*, (World Scientific, Singapore) 1984.
- ²⁰ A. Yazdani and A. Kapitulnik, Phys. Rev. Lett. **74**, 3037 (1995).
- ²¹ K.-H. Wagenblast, A. van Otterlo, R. Fazio, and G. Schön, (unpublished)
- ²² N. Trivedi, R. Scalettar and M. Randeira, cond-mat/9509029.

TABLES

Method	ϵ -exp ^(a)	(1/N)-exp ^(b)	MC ^(c)
$\frac{h}{4e^2}\sigma^* =$	0.315	0.251	0.285

TABLE I. The value of the universal conductance obtained by various methods are reported for comparison. a) this work, b) - c) Ref. 7